

Talk to your doctor about lowering your LDL-C (bad cholesterol)



In order to determine your recommended LDL-C level, your doctor needs to know about your condition, progress, and goals. This quick guide helps you gather all of that information in one place, so you can be prepared to have a productive conversation about the appropriate treatment plan for you.

Just fill out the guide, download it, and then email it to yourself.

1 What are your main concerns with having high LDL-C?

CHOOSE AS MANY AS YOU'D LIKE

- You've had a cardiovascular (CV) event
- You have a family history of heart disease
- You'd like to avoid having a heart attack
- You have other CV risk factors like smoking, diabetes, high blood pressure, etc.
- My loved ones are concerned
- I'm not happy with my current progress

2 When was your last LDL-C test?

3 What is your most recent LDL-C number?

4 What is your recommended LDL-C number?

For those with cardiovascular disease or diabetes with at least one additional risk factor, medical societies recommend an LDL-C level of less than 70 mg/dL. And depending on your risk factors, it could be lower. Talk to your doctor about what your recommended level is.

5 How have you tried to lower your bad cholesterol?

- Eating healthier
- Increasing physical activity
- Taking cholesterol-lowering medications
- Other lifestyle changes like quitting smoking

6 If you've taken LDL-C-lowering treatments and/or statins before, which ones have you tried?

7 Would you like to learn how Repatha can help bring down your LDL-C?

- Yes No

Stay informed. Sign up at www.repatha.com/enrollment to learn more

Amgen cannot make any clinical recommendations or render medical or nursing advice, treatment, opinions or care. This discussion guide is not intended to be a source of medical advice or treatment and does not replace in any way your doctor's independent medical advice regarding your diagnosis or treatment.

Please bring this guide with you next time you see your doctor and ask them if Repatha® is right for you.

Your answers to these questions can go a long way towards helping your doctor help you lower your LDL-C number.

1 Fill out this PDF

2 Download it

3 Email it to yourself so you have it when you go to the doctor

APPROVED USE

What is Repatha®?

Repatha® is an injectable prescription medicine used:

- to reduce the risk of major adverse cardiovascular (CV) events, such as death from cardiovascular disease, heart attack, stroke, certain types of chest pain conditions (unstable angina) requiring hospitalization, or certain types of heart surgery, in adults at increased risk for these events.

IMPORTANT SAFETY INFORMATION

Do not use Repatha® if you are allergic to evolocumab or to any of the ingredients in Repatha®.

Please see additional Important Safety Information on the following page.



IMPORTANT SAFETY INFORMATION

Do not use Repatha® if you are allergic to evolocumab or to any of the ingredients in Repatha®.

Before you start using Repatha®, tell your healthcare provider about all your medical conditions, including if you are allergic to rubber or latex, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. Repatha® is available as prefilled single-dose SureClick® autoinjectors and prefilled single-dose syringes that either contain dry natural rubber (a derivative of latex) in the needle cover or are not made with natural rubber latex. The carton and "Instructions for Use" will state if your prefilled single-dose SureClick® autoinjector or prefilled single-dose syringe contains dry natural rubber. The single-dose Pushtronex® system (on-body infusor with prefilled cartridge) is not made with natural rubber latex.

Tell your healthcare provider or pharmacist about any prescription and over-the-counter medicines, vitamins, or herbal supplements you take.

What are the possible side effects of Repatha®?

Repatha® can cause serious side effects including serious allergic reactions. Stop taking Repatha® and call your healthcare provider or seek emergency help right away if you have any of these symptoms: trouble breathing or swallowing, raised bumps (hives), rash or itching, swelling of the face, lips, tongue, throat or arms.

The most common side effects of Repatha® include: runny nose, sore throat, symptoms of the common cold, flu or flu-like symptoms, back pain, high blood sugar levels (diabetes) and redness, pain, or bruising at the injection site.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of Repatha®. Ask your healthcare provider or pharmacist for more information. Call your healthcare provider for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

APPROVED USE

What is Repatha®?

Repatha® is an injectable prescription medicine used:

- to reduce the risk of major adverse cardiovascular (CV) events, such as death from cardiovascular disease, heart attack, stroke, certain types of chest pain conditions (unstable angina) requiring hospitalization, or certain types of heart surgery, in adults at increased risk for these events.
- Along with diet and exercise to reduce low-density lipoprotein (LDL) or bad cholesterol in adults with high blood cholesterol levels called hypercholesterolemia and in adults with a type of high cholesterol called heterozygous familial hypercholesterolemia (HeFH).

Please see full Prescribing Information, including Patient Product Information.