



## Tracking my bad cholesterol (LDL)

Use this chart to keep track of your bad cholesterol number (measured in mg/dL) as you go forward on Repatha®. When you see your doctor, don't forget to **schedule a follow-up** to check your levels.

### MY GOAL:

(set by my doctor)

CHECK IN/VISIT	DATE		
<b>Example:</b> <i>Today's visit</i>	<i>3/26/2020</i>	<b>LDL</b> <i>148</i>	<b>NOTES/THOUGHTS</b> <i>Received Repatha® prescription</i>
<b>Today's visit</b>			
<b>Follow-up 1</b>			
<b>Follow-up 2</b>			
<b>Follow-up 3</b>			
<b>Follow-up 4</b>			

**Remember that if you stop taking Repatha®, your bad cholesterol can increase. You should not stop taking Repatha® without talking to your doctor.**

## Approved Use

Repatha® is an injectable prescription medicine used:

- in adults with cardiovascular disease to reduce the risk of heart attack, stroke, and certain types of heart surgery.
- along with diet alone or together with other cholesterol-lowering medicines in adults with high blood cholesterol levels called primary hyperlipidemia (including a type of high cholesterol called heterozygous familial hypercholesterolemia [HeFH]) to reduce low density lipoprotein (LDL) or bad cholesterol.

## Important Safety Information

**Do not use Repatha® if you** are allergic to evolocumab or to any of the ingredients in Repatha®.

**Please see additional Important Safety Information on page 2.**

### Important Safety Information (Continued)

**Before you start using Repatha® (evolocumab),** tell your healthcare provider about all your medical conditions, including if you are allergic to rubber or latex, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. The needle covers on the single-dose prefilled syringes and the inside of the needle caps on the single-dose prefilled SureClick® autoinjectors contain dry natural rubber. The single-dose Pushtronex® system (on-body infusor with prefilled cartridge) is not made with natural rubber latex.

Tell your healthcare provider or pharmacist about any prescription and over-the-counter medicines, vitamins, or herbal supplements you take.

### What are the possible side effects of Repatha®?

Repatha® can cause serious side effects including serious allergic reactions. Stop taking Repatha® and call your healthcare provider or seek emergency help right away if you have any of these symptoms: trouble breathing or swallowing, raised bumps (hives), rash or itching, swelling of the face, lips, tongue, throat or arms.

The most common side effects of Repatha® include: runny nose, sore throat, symptoms of the common cold, flu or flu-like symptoms, back pain, high blood sugar levels (diabetes) and redness, pain, or bruising at the injection site.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of Repatha®. Ask your healthcare provider or pharmacist for more information. Call your healthcare provider for medical advice about side effects.

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.**

**Please see Repatha® full Prescribing Information.**